Domain: Maternal Outcomes

Delivery Route

[Select one]

- Vaginal non-operative
- Vaginal operative
- Pre-labor cesarean / cesarean without labor
- Intrapartum cesarean
 - If cesarean, indication for cesarean (Tier 2) [Select multiple]
 - COVID-19
 - Maternal indication
 - Fetal indication
 - Other

Onset of Labor (Tier 2)

[Select one]

- No labor
- Spontaneous
- Spontaneous, augmented
- Induced
 - Primary indication for induction
 - COVID-19 infection
 - Hypertension / preeclampsia
 - IUGR
 - Non-reassuring fetal status
 - PROM
 - IUFD/ Stillbirth
 - Congenital malformations / fetal anomaly
 - Oligohydramnios
 - Abruption
 - Chorioamnionitis
 - Elective
 - Cholestasis
 - Polyhydramnios
 - Diabetes / Macrosomia
 - Other maternal medical complications
 - Post term (>=41 weeks)

Maternal Complications

[Select all that apply]

- Cerebral venous sinus thrombosis
- Renal failure requiring dialysis
- Arterial thrombosis cerebrovascular accident (CVA)
- Cardiomyopathy
- Pulmonary embolism
- Deep vein thrombosis
- Encephalopathy
- Disseminated intravascular coagulation (DIC)
- Myocardial infarction
- Superficial incisional surgical site infection
- Deep incisional surgical site infection
- Multisystem inflammatory syndrome (<18 years of age)
- HELLP syndrome
- Eclampsia
- Severe hypertension (BP>= 160/110) with acute administration of anti-hypertensive medication
- Pulmonary edema
- Hepatic rupture
- Impaired liver function (liver enzymes 2X ULN)
- Renal insufficiency (serum creatine > 1.1 mg/dL or doubling)
- Thrombocytopenia (platelets <100,000)
- Sepsis (infection with organ dysfunction)
- Bacteremia
- Endometritis requiring IV antibiotic therapy for > 24 hours
- Pelvic abscess.

Death

[After delivery discharge]

- Uterine packing
- Intrauterine balloon tamponade
- Uterine artery ligation
- Uterine compression sutures
- Laparotomy
- Evacuation of hematoma
- Arterial embolization
- Uterine evacuation
- Hysterectomy
- Death
- Other (specify) ______