

## Data Collection Worksheet

**Please Note:** The Data Collection Worksheet (DCW) is a tool to aid integration of a PhenX protocol into a study. The PhenX DCW is not designed to be a data collection instrument. Investigators will need to decide the best way to collect data for the PhenX protocol in their study. Variables captured in the DCW, along with variable names and unique PhenX variable identifiers, are included in the PhenX Data Dictionary (DD) files.

### PCL-C

**INSTRUCTIONS:** Below is a list of problems and complaints that people sometimes have in response to stressful life experiences. Please read each one carefully then circle one of the numbers to the right to indicate how much you have been bothered by that problem in the past month.

		Not at all	A little bit	Moderately	Quite a bit	Extremely
1.	Repeated, disturbing <i>memories, thoughts, or images</i> of a stressful experience from the past?	1	2	3	4	5
2.	Repeated, disturbing <i>dreams</i> of a stressful experience from the past?	1	2	3	4	5
3.	Suddenly <i>acting or feeling</i> as if a stressful experience were <i>happening again</i> (as if you were reliving it)?	1	2	3	4	5
4.	Feeling <i>very upset</i> when <i>something reminded you</i> of a stressful experience from the past?	1	2	3	4	5

5.	Having <i>physical reactions</i> (e.g., heart pounding, trouble breathing, sweating) when <i>something reminded</i> you of a stressful experience from the past?	1	2	3	4	5
6.	Avoiding <i>thinking about</i> or <i>talking about</i> a stressful experience from the past or avoiding <i>having feelings</i> related to it?	1	2	3	4	5
7.	Avoiding <i>activities</i> or <i>situations</i> because <i>they reminded</i> you of a stressful experience from the past?	1	2	3	4	5
8.	Trouble <i>remembering important parts</i> of a stressful experience from the past?	1	2	3	4	5
9.	<i>Loss of interest</i> in activities that you used to enjoy?	1	2	3	4	5
10.	Feeling <i>distant</i> or <i>cut off</i> from other people?	1	2	3	4	5
11.	Feeling <i>emotionally numb</i> or being unable to have loving feelings for those close to you?	1	2	3	4	5
12.	Feeling as if your <i>future</i> will somehow be <i>cut short</i> ?	1	2	3	4	5
13.	Trouble <i>falling</i> or <i>staying asleep</i> ?	1	2	3	4	5

14.	Feeling <i>irritable</i> or having <i>angry outbursts</i> ?	1	2	3	4	5
15.	Having <i>difficulty concentrating</i> ?	1	2	3	4	5
16.	Being " <i>super-alert</i> " or watchful or on guard?	1	2	3	4	5
17.	Feeling <i>jumpy</i> or easily startled?	1	2	3	4	5

**Scoring:**

Scoring instructions are adapted from the handout "Using the PTSD Checklist for DSM-IV (PCL)" available from the National Center for PTSD at [www.ptsd.va.gov](http://www.ptsd.va.gov).

A total symptom severity score (range = 17-85) can be obtained by summing the scores from each of the 17 items that have response options ranging from 1 "Not at all" to 5 "Extremely."

A provisional diagnosis can be established in three ways:

- 1) Determine whether an individual meets DSM-IV symptom criteria as defined by at least 1 B item (questions 1-5), 3 C items (questions 6-12), and at least 2 D items (questions 13-17). Symptoms that are rated as "Moderately" or above (responses 3 through 5 on individual items) are counted as present.
- 2) Determine whether the total severity score exceeds a relevant normative threshold (e.g., for primary care settings).
- 3) Combine methods (1) and (2) to ensure that an individual meets both the symptom pattern and severity threshold.

Protocol source: <https://www.phenxtoolkit.org/protocols/view/122001>