



Data Collection Worksheet

Please Note: The Data Collection Worksheet (DCW) is a tool to aid integration of a PhenX protocol into a study. The PhenX DCW is not designed to be a data collection instrument. Investigators will need to decide the best way to collect data for the PhenX protocol in their study. Variables captured in the DCW, along with variable names and unique PhenX variable identifiers, are included in the PhenX Data Dictionary (DD) files.

SUICIDE INTENT SCALE

(For Attempters)

Instructions: Interview the patient about his/her most recent suicide attempt to enable a reliable coding of the objective circumstances of the attempt, and the patient's perceptions of the attempt. The total score is computed by summing items 1-15. If an item is "Not applicable" code it "-8." Do not include "-8's" when calculating the total score.

I. Objective Circumstances Related to Suicide Attempt

1. Isolation

- 0 [] Somebody present
- 1 [] Somebody nearby, or in visual or vocal contact
- 2 [] No one nearby or in visual or vocal contact

2. Timing

- 0 [] Intervention is probable
- 1 [] Intervention is not likely
- 2 [] Intervention is highly unlikely

3. Precautions against Discovery/Intervention

- 0 [] No precautions
- 1 [] Passive precautions (avoiding others but doing nothing to prevent intervention; alone in room with unlocked door)
- 2 [] Active precautions (locked door)

4. Acting to Get Help During/After Attempt

- 0 Notified potential helper regarding attempt
- 1 Contacted but did not specifically notify potential helper regarding attempt
- 2 Did not contact or notify potential helper

5. Final Acts in Anticipation of Death (e.g., will, gifts, insurance)

- 0 None
- 1 Thought about or made some arrangements
- 2 Made definite plans or completed arrangements

6. Active Preparation for Attempt

- 0 None
- 1 Minimal to moderate
- 2 Extensive

7. Suicide Note

- 0 Absence of note
- 1 Note written, but torn up; note thought about
- 2 Presence of note

8. Overt Communication of Intent Before the Attempt

- 0 None
- 1 Equivocal communication
- 2 Unequivocal communication

II. Self-Report

9. Alleged Purpose of Attempt

- 0 To manipulate environment, get attention, revenge
- 1 Components of "0" and "2"
- 2 To escape, surcease, solve problems

10. Expectations of Fatality

- 0 [] Thought that death was unlikely
- 1 [] Thought that death was possible but not probable
- 2 [] Thought that death was probable or certain

11. Conception of Method's Lethality

- 0 [] Did less to self than he thought would be lethal
- 1 [] Wasn't sure if what he did would be lethal
- 2 [] Equaled or exceeded what he thought would be lethal

12. Seriousness of Attempt

- 0 [] Did not seriously attempt to end life
- 1 [] Uncertain about seriousness to end life
- 2 [] Seriously attempted to end life

13. Attitude Toward Living/Dying

- 0 [] Did not want to die
- 1 [] Components of "0" and "2"
- 2 [] Wanted to die

14. Conception of Medical Rescuability

- 0 [] Thought that death would be unlikely if he received medical attention
- 1 [] Was uncertain whether death could be averted by medical attention
- 2 [] Was certain of death even if he received medical attention

15. Degree of Premeditation

- 0 [] None; Impulsive
- 1 [] Suicide contemplated for three hours or less prior to attempt
- 2 [] Suicide contemplated for more than three hours prior to attempt

III. Other Aspects (Not Included in Total Score)

16. Reaction to Attempt

- 0 [] Sorry about attempt; feels foolish, ashamed (circle which one)

1 Accepts both attempt and its failure

2 Regrets failure of attempt

17. Visualization of Death

0 Life-after-death, reunion with decedents

1 Never ending sleep, darkness, end-of-things

2 No conceptions of, or thoughts about death

18. Number of Previous Attempts

0 None

1 One or two

2 Three or more

19. Relationship between Alcohol Intake and Attempt

0 Some alcohol intake prior to but not related to attempt, reportedly not enough to impair judgment, reality testing

1 Enough alcohol intake to impair judgment, reality testing and diminish responsibility/impulse control

2 Intentional intake of alcohol in order to facilitate implementation of suicide attempt

20. Relationship between Drug Intake and Attempt (narcotics, hallucinogens, etc. drug is not the method used to suicide)

0 Some drug intake prior to but not related to attempt, reportedly not enough to impair judgment, reality testing

1 Enough drug intake to impair judgment, identity testing and diminish responsibility/impulse control

2 Intentional drug intake in order to facilitate implementation of suicide attempt

Scoring:

Each of the 15 items is scored from 0 to 2 and added together to give a total score from 0 to 30.

Protocol source: <https://www.phenxtoolkit.org/protocols/view/640301>